APPENDIX A



Your Discoveries Begin with US.

Search: - Choose a product line -

Home | Ordering Info | Technical Help | About ATCC | Contact Us

Search

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's <u>Material Transfer Agreement</u>.

Customers in Europe, Australia, Japan, Korea, New Zealand and Talwan must contact a <u>local distributor</u> for pricing information and to place an order for ATCC cultures and products.

Cell Line	S		and the second s			
ATCC Number	Order this item	Price:	\$215.00			
Designation	W6/32	Depositors:	P Parham			
Biosafety Le	vel: 1 Shipped: frozen	Isotype:	IgG2a			
Medium & Serum:	See Propagation	Growth Properties:	suspension			
Organism:	Mus musculus (B cell); Mus musculus (myeloma) (mouse (B cell); mouse (myeloma))	Morphology:	lymphoblast			
Tissue:	B lymphocyte; hybridoma	B lymphocyte; hybridoma				
Cellular Products: Permits/Form	Immunoglobulin; monoclonal antibo ns: In addition to the MTA mentioned a be required for the transfer of this a ultimately responsible for obtaining	bove, other ATCC an ATCC material. Anyon	d/or regulatory permits may ne purchasing ATCC material is			
ultimately responsible for obtaining the permits. Please <u>click here</u> for information regarding the specific requirements for shipment to your location. Related Cell Culture Products						
Comments:	Animals were immunized with cells from human tonsils. Spleen cells were fused with P3X63Ag8 myeloma cells. The antibody reacts with monomorphic determinants on HLA A,B,C molecules. The antibody is cytotoxic. The antibody cross-reacts with ape, monkey and cow cells. It does not react with rabbit cells. Tested and found negative for extremella virus (mousepox).					
Propagation:	ATCC medium: Dulbecco's modified Ea bovine serum, 10%.	CC medium: Dulbecco's modified Faule's medium with 4.5 g/l, glusses 2001, Guide				
Subculturing:	Cultures can be maintained by addition or replacement of fresh medium. Start cultures at 2 X 10 exp5 cells/ml and maintain between 1 X 10 exp5 and 1 X 10 exp6 cells/ml.					
Fluid Renewal:	Every 2 to 3 days					
References:	955: Brodsky FM, Parham P. Monomorphic anti-HLA-A,B,C monocional antibodies detecting molecular subunits and combinatorial determinants. J. Immunol. 128: 129-135, 1982. PubMed: 6172474 956: Barnstabl CJ, et al. Production of monocional antibodies to group A erythrocytes, HLA and other human cell surface antigens new tools for genetic analysis. Cell 14:					

9-20, 1978. PubMed: 667938
22830: Parham P, et al. Use of a monoclonal antibody (W6/32) in structural studies of HLA- A,B,C, antigens. J. Immunol. 123: 342-349, 1979. PubMed: 87477
32767: Solomon KR, et al. The association between glycoslphosphatidylinositol-anchored proteins and hetrotrimeric G protein alpha subunits in lymphocytes. Proc. Natl. Acad. Sci. USA 93: 6053-6058, 1996. PubMed: 8650218

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your word search in Cell Lines by typing in your search criteria below or returning to the Cell Lines monu. To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for AND), | (for QR), ^ (for AND NOT). An asterisk (*) is used as the wildcard. For more information please review the Search Help.

w6/32 Word Search Clear Search

For query options, please read the search help.

Home Page Archive

Home Ordering Info Technical Help About ATCC Contact Us Privacy Policy Terms of Use ATCC MTA

© 2003 American Type Culture Collection (ATCC).
All rights reserved.



Your Discoveries Begin with US.

Search: - Choose a product line - _

Home Ordering Info | Technical Help | About ATCC | Contact Us

Search

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's <u>Material Transfer Agreement</u>.

Customers in Europe, Australia, Japan, Korea, New Zealand and Taiwan must contact a $\frac{\log d}{d}$ distributor for pricing information and to place an order for ATCC cultures and products.

Cell Lines	3				İ		
ATCC Number	r: HB-54	Order	this item	Price:	\$283.00		
Designation:	MA2.1			Depositors:	P Parham		
Biosafety Lev	<u>rel</u> : 1	Shipped:	frozen	Isotype:	lgG1		
Medium & Scrum:	See Propagation			Growth Properties:	suspension		
Organism:	musculus	<i>culus</i> (B cell) (myeloma) use (myelom	(mouse (B	Morphology:	lymphoblast		
Tissue:	B lympho	B lymphocyte; hybridoma					
Cellular Products: Immunoglobulin; monoclonal antibody; against HLA A2, B17 Permits/Forms: In addition to the MTA mentioned above, other ATCC and/or regulatory permits may be required for the transfer of this ATCC material. Anyone purchasing ATCC material is ultimately responsible for obtaining the permits. Please click here for Information regarding the specific requirements for shipment to your location.							
	Related Cell Culture Products						
Comments:	Spicen cells Mutation at complete lo	unimals were immunized with papaln solubilized HLA A2 and B17 antigens from the J-Y ell line. Spleen cells were fused with NS-1 myeloma cells. Sutation at positions 62 to 66 of the alpha 1 domain of the HLA-A2 molecule results in complete loss of binding of MA2.1. [39935] ested and found negative for extromelia virus (mousepox).					
Propagation:		ATCC medium: RPMI 1640 medium, 80%; fetal bovine serum, 20%					
	Subculturing: Cultures can be maintained by addition or replacement of fresh medium. Start cultures at 1 X 10 exp5 cells/ml and maintain between 1 X 10 exp5 and 1 X 10 exp6 cells/ml.						
Fluid Renewal:		very 2 to 3 days					
References:	930: McMichael AJ, et al. A monocional antibody that recognizes an antigenic determinant shared by HLA A2 and B17. Hum. Immunol. 1: 121-129, 1980. PubMed: 6167544 39935: Santos-Aguado J, et al. Molecular characterization of serologic recognition sites in the human HLA-A2 molecule. J. Immunol. 141: 2811-2818, 1988. PubMed: 2459229 39936: Hogan KT, Brown SL. Localization and characterization of serologic epitopes on HLA-A2. Hum. Immunol. 33: 185-192, 1992. PubMed: 1377666						

Notices and Disclaimers

ATCC products are intended for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as tost tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your word search in Cell Lines by typing in your search criteria below or returning to the <u>Cell Lines menu</u>. To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for AND), I (for OR), ^ (for AND NOT). An asterisk (*) is used as the wildcard. For more information please review the <u>Search Help</u>.

hb-54 Word Search Clear Search

For query options, please read the search help.

Home Page Archive

Home Ordering Info Technical Help About ATCC Contact Us Privacy Policy Terms of Use ATCC MTA

© 2003 American Type Culture Collection (ATCC).
All rights reserved.



Your **Discoveries**Begin with US.

Search: - Choose a product line - ____

Home | Ordering Info | Technical Help | About ATCC | Contact Us

Search

Before submitting an order you will be asked to read and accept the terms and conditions of ATCC's <u>Material Transfer Agreement</u>.

Customers In Europe, Australia, Japan, Korea, New Zealand and Talwan must contact a $\underline{local\ distributor}$ for pricing information and to place an order for ATCC cultures and products.

Cell Lines		,				
ATCC Number:	HB-149 Order this item	Price:	\$283.00			
Designation:	L368	Depositors:	L Lampson			
Biosafety Leve	el: 1 Shipped: frozen	Isotype:	IgG1; kappa light chain			
Medium & Serum:	Sec Propagation	Growth Properties:	suspension			
Organism:	Mus musculus (B cell); Mus musculus (myeloma) (mouse (B cell); mouse (myeloma))	Morphology:	lymphoblast			
Tissue:	B lymphocyte; hybridoma					
Cellular Products:	immunoglobulin; monoclonal antibody; against human beta-2-microglobulin (beta 2 microglobulin)					
Permits/Form	s: In addition to the MTA mentioned ab- be required for the transfer of this AT ultimately responsible for obtaining t regarding the specific requirements f	FCC material, Anyo he permits. Picase	ne purchasing ATCC material is click here for information			
		<u>Reli</u>	ated Cell Culture Products			
	Animals were immunized with the RPMI 8866 human B cell line. Spiecn cells were fused with NS-1 myeloma cells. The antibody does not cross-react with mouse, rat, cat or monkey beta-2-microglobulin. The antibody is not cytotoxic. Tested and found negative for ectromolla virus (mousepox).					
Propagation:	ATCC medium: Dulbecco's modified Eagle's medium with 4.5 g/L glucose, 0.1 mM hypoxanthine and 0.016 mM thymidine, 90%; fetal bovine serum, 10%					
	cultures can be maintained by addition or replacement of fresh medium. Start cultures at 2 X 10 exp5 cells/ml and maintain between 1 X 10 exp5 and 1 X 10 exp6 cells/ml.					
Fluid Renewal:	Every 3 to 4 days					
References:	1000: Lampson LA, et al. Striking paucity of HLA-A, B, C and beta2-microglobulin on human neuroblastoma cell lines. J. Immunol. 130: 2471-2478, 1983. PubMed: 6187860					

Notices and Disclaimers

ATCC products are intend id for laboratory research purposes only. They are not intended for use in humans.

While ATCC uses reasonable efforts to include accurate and up-to-date information on this site, ATCC makes no warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. ATCC does not warrant that such information has been confirmed to be accurate.

All prices are listed in U.S. dollars and are subject to change without notice. A discount off the current list price will be applied to most cultures for nonprofit institutions in the United States and Canada. Cultures that are ordered as test tubes or flasks will carry an additional laboratory fee. Fees for permits, shipping, and handling may apply.

You may continue your word search in Cell Lines by typing in your search criteria below or returning to the Cell Lines menu. To search another product line, choose one from the dropdown box at the top. For complex searches using boolean operators, the following characters must be used: & (for AND), | (for OR), ^ (for AND NOT). An asterisk (*) is used as the wildcard. For more information please review the Search Help.

368 Word Search Clear Search

For query options, please read the search help.

Home Page Archive

Home Ordering Info Technical Help About ATCC Contact Us Privacy Policy Terms of Use ATCC MTA

© 2003 American Type Culture Collection (ATCC).
All rights reserved.